



WEST Search History

DATE: Tuesday, June 28, 2005

Hide?	<u>Set</u> <u>Name</u>	<u>Query</u>	<u>Hit</u> <u>Count</u>
	<i>DB=PGPB,USPT,USOC,EPAB,JPAB,DWPI,TDBD; PLUR=YES; OP=OR</i>		
<input type="checkbox"/>	L34	L33 and APO-3	34
<input type="checkbox"/>	L33	L32 and apoptosis	172
<input type="checkbox"/>	L32	l30 and antagonist	270
<input type="checkbox"/>	L31	L30 and antaognist	0
<input type="checkbox"/>	L30	(APO)same(antibod?)	952
<input type="checkbox"/>	L29	L28 and apo-3	2
<input type="checkbox"/>	L28	424/178.1.ccls.	1051
<input type="checkbox"/>	L27	L26 and APO-3	0
<input type="checkbox"/>	L26	424/142.1.ccls.	241
<input type="checkbox"/>	L25	L24 and Apo-3	1
<input type="checkbox"/>	L24	424/133.1.ccls.	644
<input type="checkbox"/>	L23	L22 and APO-3	3
<input type="checkbox"/>	L22	424/130.1.ccls.	1634
<input type="checkbox"/>	L21	L18 and APO-3	0
<input type="checkbox"/>	L20	L18 and Apo-3	0
<input type="checkbox"/>	L19	L18 and Apo-3 antibody	217182
<input type="checkbox"/>	L18	435/7.21.ccls.	2559
<input type="checkbox"/>	L17	435/7.21,ccls.	0
<input type="checkbox"/>	L16	L15 and Apo-3 receptor	209375
<input type="checkbox"/>	L15	L14 and activation	47
<input type="checkbox"/>	L14	L13 and apoptosis	47
<input type="checkbox"/>	L13	L12 and inhibit	47
<input type="checkbox"/>	L12	L11 and antibod?	47
<input type="checkbox"/>	L11	L10 and Apo-3	47
<input type="checkbox"/>	L10	435/7.1.ccls.	10005
<input type="checkbox"/>	L9	L8 and anti-Apo-3	9
<input type="checkbox"/>	L8	L7 and APO-3	157
<input type="checkbox"/>	L7	(ashkenazi)	2569
<input type="checkbox"/>	L6	(Avi)adj(J)adj(ashkenazi)	0
<input type="checkbox"/>	L5	(A)adj(ashkenzai)	0

<input type="checkbox"/>	L4	(ashkenzai)same(APO-3)	0
		<i>DB=USPT; PLUR=YES; OP=OR</i>	
<input type="checkbox"/>	L3	US-6469144-B1.did.	1
		<i>DB=PGPB,USPT,USOC,EPAB,JPAB,DWPI,TDBD; PLUR=YES; OP=OR</i>	
<input type="checkbox"/>	L2	(APO-3)same(antibod?)	19
<input type="checkbox"/>	L1	(APO-3)same(antibod?)same(antagonist)same(apoptosis)same(NF)adj (kappa)adj(B)adj(activation)	0

END OF SEARCH HISTORY



All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Book

Search PubMed for [] Go Clear

Limits Preview/Index History Clipboard Details

Display Abstract Show 20 Sort by Send to

About Entrez

Text Version

All: 1 Review: 0

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

Special Queries

LinkOut

My NCBI (Cubby)

Related Resources

Order Documents

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

1: Science. 1989 Jul 21;245(4915):301-5.

Related Articles, Links

Monoclonal antibody-mediated tumor regression by induction of apoptosis.**Trauth BC, Klas C, Peters AM, Matzku S, Moller P, Falk W, Debatin KM, Krammer PH.**

Institute for Immunology and Genetics, German Cancer Research Center, Heidelberg.

To characterize cell surface molecules involved in control of growth of malignant lymphocytes, monoclonal antibodies were raised against the human B lymphoblast cell line SKW6.4. One monoclonal antibody, anti-APO-1, reacted with a 52-kilodalton antigen (APO-1) on a set of activated human lymphocytes, on malignant human lymphocyte lines, and on some patient-derived leukemic cells. Nanogram quantities of anti-APO-1 completely blocked proliferation of cells bearing APO-1 in vitro in a manner characteristic of a process called programmed cell death or apoptosis. Cell death was preceded by changes in cell morphology and fragmentation of DNA. This process was distinct from antibody- and complement-dependent cell lysis and was mediated by the antibody alone. A single intravenous injection of anti-APO-1 into nu/nu mice carrying a xenotransplant of a human B cell tumor induced regression of this tumor within a few days. Histological thin sections of the regressing tumor showed that anti-APO-1 was able to induce apoptosis in vivo. Thus, induction of apoptosis as a consequence of a signal mediated through cell surface molecules like APO-1 may be a useful therapeutic approach in treatment of malignancy.



PMID: 2787530 [PubMed - indexed for MEDLINE]

Display Abstract Show 20 Sort by Send to

[Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)

Department of Health & Human Services

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)



All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search PubMed for [] Go Clear

Limits Preview/Index History Clipboard Details

Display Abstract Show 20 Sort by Send to

About Entrez

Text Version

All: 1 Review: 0

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

Special Queries

LinkOut

My NCBI (Cubby)

Related Resources

Order Documents

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

1: Curr Biol. 1996 Dec 1;6(12):1669-76.

Related Articles, Links

Apo-3, a new member of the tumor necrosis factor receptor family, contains a death domain and activates apoptosis and NF-kappa B.

Marsters SA, Sheridan JP, Donahue CJ, Pitti RM, Gray CL, Goddard AD, Bauer KD, Ashkenazi A.

Department of Molecular Oncology, Genentech, Inc., South San Francisco, California 94080-4918, USA.

BACKGROUND: Two receptors that contain the so-called "death domain" have been described to date: tumor necrosis factor receptor 1 (TNFR1) and Fas/Apo-1 (CD95); both belong to the TNFR gene family. The death domain of TNFR1 mediates the activation of programmed cell death (apoptosis) and of the transcription factor NF-kappa B, whereas the death domain of CD95 only appears to activate apoptosis. **RESULTS:** We have identified an additional member of the TNFR family, which we have named Apo-3. Apo-3 is a transmembrane protein of approximately 47 kDa that has similarity of members of the TNFR family in its extracellular, cysteine-rich domains. In addition, Apo-3 resembles TNFR1 and CD95 in that it contains a cytoplasmic death domain. The Apo-3 gene mapped to human chromosome 1p36.3, and Apo-3 mRNA was detected in several human tissues, including spleen, thymus, peripheral blood lymphocytes, small intestine and colon. Ectopic expression of Apo-3 in HEK293 or HeLa cells induced marked apoptosis. CrmA, a poxvirus inhibitor of Ced-3-like proteases which blocks death signaling by TNFR1 and CD95, inhibited Apo-3-induced apoptosis. Ectopic expression of Apo-3 also induced the activation of NF-kappa B. Apo-3 did not specifically bind to the Apo-2 ligand, suggesting the existence of a distinct ligand for Apo-3. **CONCLUSIONS:** These results identify Apo-3 as a third member of the TNFR family that activates apoptosis, and suggest that Apo-3, TNFR1 and CD95 engage a common apoptotic cell-death machinery. Apo-3 resembles TNFR1 because it can stimulate NF-kappa B activity and regulate apoptosis. Apo-3 mRNA is expressed in various tissues, consistent with the possibility that this receptor may regulate multiple signaling functions.